

Working together to eliminate cyanide poisoning, konzo, tropical ataxic neuropathy (TAN) and
neurolethyrism



CCDNN

News

Cassava Cyanide Diseases & Neurolethyrism Network

(ISSN 1838-8817 (Print): ISSN 1838-8825 (Online))

Issue Number 21, June 2013

Contents

Editorial..... 1

***Lathyrus sativus*: To eat or not to eat?..... 1**

Konzo control and cassava safety using the popular wetting method to remove poisonous cyanogens 2

The cultivation, consumption and research progress of *Lathyrus* in China..... 4

Impact of cross-sectoral approach to addressing konzo in DRC 8

Safety and toxicity profile of grass pea (*Khesari dal*): A reappraisal in the Indian scenario 10

CCDNN Coordination:

Prof Fernand LAMBEIN, Coordinator
Ghent University, Institute for plant Biotechnology Outreach (IPBO)
Proeftuinstraat 86 N1, B-9000 Ghent, Belgium
Phone: +32 484 417 5005
E-mail: Fernand.Lambein@ugent.be or Fernand.Lambein@gmail.com

Dr. Delphin DIASOLUA NGUDI
Ghent University, Institute for plant Biotechnology Outreach (IPBO)
Proeftuinstraat 86 N1, B-9000 Ghent, Belgium
Phone: +32 92 64 52 82
Email: ddiasolu@yahoo.com

Editorial Board:

J.P. Banea, Julie Cliff, Arnaldo Cumbana, Ian Denton, D. Diasolua Ngudi, F. Lambein, J. Howard Bradbury, N.L.V.Mlingi, Humberto Muquingue, Bala Nambisan, Dulce Nhassico, S.L.N. Rao, Ivan Ingelbrecht.

Country Contacts:

Cameroon: E.E. Agbor;
D.R. Congo: D. Diasolua Ngudi and J. Nsimire Chabwine; *Indonesia*: A. Hidayat;
Mozambique: Anabela Zacarias;
Nigeria: M.N. Adindu and P.N. Okafor

Website: www.anu.edu.au/BoZo/CCDNN (temporarily)
<http://www.ugent.be/we/genetics/ipbo/en/networking/ccdnn/overview.htm>

Editorial

With CCDNN21 we start the second decade of regular bi-annual publication. After a smooth transition in 2012, the newsletter is now published by an international editorial committee coordinated from Ghent University. With this issue we further extend the geographical coverage with contributions from China and India on aspects of neurolethyrism. A group from Australia and one from the US report on aspects of konzo in central Africa. These results support our hope that perhaps in another decade, konzo and neurolethyrism will disappear and only have historical importance. As a way to consolidate recent progress, an international “Cassava Cyanide Diseases Workshop” will be convened shortly, probably in Kinshasa, D.R. Congo. Detailed notifications will follow. We start with the abstract of a lecture presented at the 1st Legume Society Conference, 9-10 May 2013 in Novi Sad, Serbia.

***Lathyrus sativus*: To eat or not to eat?**

In its long history as a cultivated crop, grass pea (*Lathyrus sativus* L.) has received praise and blame. In the Pharaonic era, grass pea was part of funeral offerings found in the pyramids. In drought prone areas of Ethiopia and the Indian Sub-continent, grass pea is considered a life-insurance

crop. Already in antiquity, the link was made between overconsumption of grass pea seeds and a crippling neurological disorder, later coined *neurolathyrism*. Overemphasis on suspected toxicity of the seeds has led to neglect of the exceptional positive agronomic properties of the plant. In normal situations, *neurolathyrism* is virtually non-existent. The etiology of *neurolathyrism* had been oversimplified as being caused by a single metabolite and factors such as oxidative stress, mineral content, the total diet, emotional stress and deficiency of essential amino acids were neglected. Recent epidemiological and pharmacological studies have indicated the importance of essential amino acids and oxidative stress in the incidence of *neurolathyrism*. The same metabolite that is considered the cause of *neurolathyrism* is also present in Ginseng and patented in China as a haemostatic drug.

The drought tolerant grass pea is the survival food for the poor during drought-triggered famines. It is the most efficient nitrogen fixer among commercial legumes, grows on poor soil and yields even better in moderate salinity. Grass pea can become a wonder crop if it can shed the double stigma of being a toxic plant and the food for the poorest of the poor. All strategies including genetic transformation need to be exploited.

Fernand LAMBEIN and Yu-Haey KUO
IPBO, Ghent University, K.L. Ledeganckstraat 35,
B-9000 Ghent, Belgium

Konzo control and cassava safety using the popular wetting method to remove poisonous cyanogens

Konzo is an irreversible paralysis of the legs that occurs mainly in children and young women after childbirth, associated with (1) a high intake of cyanogens from a monotonous diet of bitter cassava and (2) a diet that is deficient in protein, in particular the S-containing amino acids methionine and cysteine/cystine, that are needed to detoxify cyanide to produce thiocyanate.¹⁻³ Konzo is a non-

progressive upper motor neuron disease of sudden onset that was first discovered by Trolli⁴ in 1938 in Popokabaka Health Zone, Bandundu Province, Belgian Congo, now the Democratic Republic of Congo (DRC). A recent estimate of the number of konzo cases in the DRC was 100,000. Konzo also occurs in lesser amount in Mozambique, Tanzania, Cameroon, Central African Republic and was recently reported from

Angola.⁵ It has been controlled for the first time ever, in the same Health Zone in which it was first discovered, by reducing greatly the intake of cyanogens from cassava flour.⁶ The wetting method has now been shown to be a popular method to ensure food safety.⁷

The cyanogen content of cassava flour, prepared by traditional methods of soaking peeled roots in water or sun drying, is greatly reduced using the simple wetting method, as follows.⁸⁻¹⁰ The flour is added to a bowl and the level marked on the inside of the bowl. Water is added with mixing until the level of the wet flour comes up to the mark. The wet flour is spread out on a mat in a thin layer and the enzyme (linamarase) breaks down the cyanogen (linamarin) to acetone cyanohydrin that hydrolyses spontaneously at about pH 6.5, liberating hydrogen cyanide (HCN) gas, which escapes from the sticky wet flour. The process requires about 2 hours in the sun or 5 hours in the shade for nearly complete removal of cyanogens.

The women in the village of Kay Kalenge were educated about the poison present in cassava flour and shown how to use the wetting method to remove it. They accepted the method and used it on their cassava flour. Monthly visits were made to the village by Sister Mandombi's Caritas team to help the women use the method correctly and every four months Dr Banea's full team visited the village to check on any new cases of konzo and measure the total cyanide content of flour samples just before they were used to prepare the thick porridge (fufu), which is eaten with something to give it flavour such as pounded, boiled cassava leaves.¹¹ Analysis of urine samples from school children for thiocyanate, produced by detoxification

of ingested cyanide, gave a good measure of cyanide intake over previous days.

No new cases of konzo occurred in Kay Kalenge during the 18 month study, the total cyanide content of cassava flour samples fell to below the WHO safe level of 10 ppm and the urinary thiocyanate content of school children reduced to safe levels.⁶ In 2011-2 we made a 12 month intervention in three villages in Boko Health Zone where there were 61 konzo cases and a mean prevalence rate of 4.6%. The intervention has proved to be just as successful as that in Kay Kalenge with no new cases of konzo occurring in the villages and large reductions in cassava flour cyanide and in urinary thiocyanate levels. Also the people of surrounding villages were willing to use the wetting method.¹² Another successful intervention in three more high prevalence konzo villages is nearing completion also with AusAID support.

We returned to Kay Kalenge village 14 months after the intervention ceased. We were pleased to find that there were no new cases of konzo in the village, all the women were still using the wetting method and the total cyanide content of flour samples and urinary thiocyanate levels were low. But most pleasing was the fact that the wetting method had spread by word of mouth to three nearby villages.⁷

Why is the wetting method so popular with rural women? Because it is simple and does not require much extra work or additional equipment, except for a basin and a mat. The treated flour produces high quality good tasting fufu, because of removal of the bitter taste of linamarin, and the fufu can be stored for up to three days.^{6,13} Most importantly, it ensures food safety. It has been shown that high cyanide intake from cassava at levels not sufficient to cause konzo, is associated with impaired intellectual development in children.¹⁴ To avoid this undesirable effect on their children, many families in tropical Africa and elsewhere who consume cassava flour may wish to use the wetting method to ensure food safety.

We gratefully acknowledge the financial support of

the Australian Agency for International Development (AusAID).

References

- ¹ Ministry of Health Mozambique, 1984. Mantakassa; an epidemic of spastic paraparesis associated with chronic cyanide intoxication in a cassava staple area in Mozambique. 1. Epidemiology and clinical laboratory findings in patients. Bull. WHO, 62: 477-484.
- ²Cliff, J., Martensson, J., Lundquist, P., Rosling H., Sorbo, B., 1985. Association of high cyanide and low sulphur intake in cassava induced spastic paraparesis. Lancet 11, 1211-1213.
- ³ Howlett, W.P., Brubaker, G.R., Mlingi, N., Rosling, H., 1990. Konzo, an epidemic upper motor neuron disease studied in Tanzania. Brain 113, 223-235.
- ⁴ Trolli, G., 1938. Paraplégie spastique épidémique, "Konzo" des indigènes du Kwango. In: Trolli, G., ed. Résumé des observations réunies, au Kwango, au sujet de deux affections d'origine indéterminée. Brussels: Fonds reine Elisabeth, 1-36.
- ⁵ Bettencourt, M.S., Paquise, M.M., Zangulo, A., Reseande, I. 2011. Tropical spastic paraparesis (konzo): a major neurologic problem in Caungula-Angola: first report in Angola. CCDN News, No 18, P 4.
- ⁶ Banea, M., Nahimana, G., Mandombi, C., Bradbury, J.H., Denton, I.C., Kuwa, N. 2012. Control of konzo in DRC using the wetting method on cassava flour. Food Chem. Toxicol. 50, 1517-1523.
- ⁷ Banea, J.P., Bradbury, J.H., Mandombi, C., Nahimana, D., Denton, I.C., Kuwa, N., Tshala Katumbay, D. 2013. Control of konzo in the DRC by removal of cyanogens from cassava flour using the wetting method. Food Nutrition Bulletin, submitted for publication.
- ⁸ Bradbury, J.H., 2006. Simple wetting method to reduce cyanogen content of cassava flour. J. Food Comp. Anal. 19, 388-393.
- ⁹ Cumbana, A., Mirione, E., Cliff, J., Bradbury, J.H., 2007. Reduction of cyanide content of cassava flour in Mozambique by the wetting method. Food Chem. 101, 894-897.
- ¹⁰ Bradbury, J.H., Denton, I.C., 2010. Rapid wetting method to reduce cyanogen content of cassava flour. Food Chem., 121, 591-594.
- ¹¹ Diasolua Ngudi, D., Kuo, Y.H., Lambein, F. 2003 Amino acid profiles and protein quality of cooked cassava leaves or sakasaka. J. Sci. Food Agric. 83, 529-534.
- ¹² Banea, J.P., Bradbury, J.H., Mandombi, C., Nahimana, D., Denton, I.C., Kuwa, N., Tshala Katumbay, D. 2013. Control of konzo by detoxification of cassava flour in three villages in the Democratic Republic of Congo. Food

Chem. Toxicol., in press.

¹³ Bradbury, J.H., Cliff, J. Denton, I.C. 2011. Uptake of wetting method in Africa to reduce cyanide poisoning and konzo from cassava. *Food Chem. Toxicol.* 49, 539-542.

¹⁴ Boivin, M.J., Okitunda, D., Bumoko, G.M., Sombo, M.T., Mumba, D., Tylleskar, T., Page, C.F., Muyembe, J.T., Tshala Katumbay, D., 2013. Neuropsychological effects of konzo: a neuromotor disease associated with poorly processed cassava. *Pediatrics* 131, April 2013, e1231-e1238.

J.H. Bradbury,¹ J.P.Banea,² C. Mandombi,³ D. Nahimana,² I.C. Denton,¹ D. Tshala Katumbay,⁴ N. Kuwa²

¹ EEG, Research School of Biology, Australian National University, Canberra, Australia.

²Programme National de Nutrition (PRONANUT), Kinshasa, DRC

³ Hopital General de Référence, Zone de Santé de Popokabaka, DRC.

⁴ Centre Neuropsychopathologique, Université de Kinshasa, DRC.

The cultivation, consumption and research progress of *Lathyrus* in China

Grass pea (*Lathyrus sativus* L.), also named horse-tooth-like pea, three-edge pea or *song liu* pea since its leaves are similar to the ones of pine and willow in China, is an annual pulse crop belonging to the family *Fabaceae* and the tribe *Vicieae*. It has been cultivated in China for over 600 years according to historic records. During the famine times grass pea used to be the only lifesaving food for both human consumption and animal feeding. At present, grass pea has been extensively cultivated and naturalized as a forage crop in fields with different soil types. Here, we mainly report on the cultivation, consumption and research progress of grass pea in China.

The cultivation history and current status of grass pea

The cultivation of *Lathyrus* species in China has a long history, the earliest historical evidence comes from the *Jiu Huang Ben Cao* (on wild food plants for use in emergency) by Su Zhu in the early Ming dynasty (about 1400s). Nowadays, the main *Lathyrus* species cultivated in China are *L. sativus*,

L. cicera, *L. odoratus*, *L. tingitanus* and *L. quinquenervius*.^{1,2} Among them, *L. sativus* has a wide range of cultivation in the northeast, northwest and southwest of China. In the 1960s, Chinese people suffered from a severe grain deficiency due to drought and many other reasons. As a result grass pea was over-consumed as an exclusive staple food, especially in the low-income and food deficit areas, which led to the occurrence of serious neurolathyrism. Therefore, grass pea was forbidden by the government to be consumed as food in the 1970s. In spite of this, grass pea is still cultivated for animal forage, especially in the northwest of China^{3,4} and Inner Mongolia.⁵ Dingxi region in Gansu province, including one administrative area and six counties, is the main cultivating area for grass pea in China, producing approximate 5000 tons of seeds in 2012,⁷ occupying 3.3% of the total forage area for its abundant nutrients,⁶ and the yield of grass seeds is about 1.13 ton per hectare.

The consumption of grass pea and neurolathyrism

In addition to the extensive use of its seeds as forage, grass pea seedlings are also utilized as a potherb.⁸ At present, Chinese people usually prepare it for consumption by baking the seeds, stir-frying or boiling soup using its sprouts mixed with other foods, like eggs, meat and shrimp or other vegetables. Recently, a research group reported that the seedlings of grass pea which is called grass pea sprouts could be eaten as either cold dishes or hot pot vegetables.⁹ We have investigated 15 large supermarkets in Lanzhou, and found 4 supermarkets selling grass pea sprouts provided by special producers, however, the amount of sales is quite small. Moreover, *L. quinquenervius* is now used as a precious herbal medicine in the northeast of China for its effectiveness in treating arthritis and headaches. A recent study has reported the value of grass pea as a green manure for improving soil nutrients.¹⁰

According to the records of Gansu Provincial Archives, neurolathyrism occurred in 8 counties in the 1970s, including Gangu, Qin'an, Longxi, Hoeryong, Tongwei, Wushan, Yuzhong and

Tianshui and was caused by excessive consumption of grass pea. A total of 2315 cases of neuropathy were identified in these areas. The main symptoms of neuropathy are dizziness, headache, lacking of energy, lassitude and nausea, further development of the disease could cause lower limb numbness and weakness, leg twitching, trembling and stiffness. In severe cases patients walk becomes difficult with scissor gait. Of all the cases, 85.6% were man, aged from 18 to 40, and 204 cases were bedridden. In the 1990s, Gansu Provincial Hospital also surveyed neuropathy in the outbreak areas which were linked with poverty and illiteracy. After neurologic examination of 2128 farmers, 119 were diagnosed to have neuropathy, accounting for a prevalence of 5.6%. The diagnosed patients consisted of 95 men and 24 women, among them 65 were hard laborers, accounting for 7.9% of the total labor forces. Clinical researches revealed that the motor neuron injury of neuropathy was related to prolonged overconsumption of grass pea, and the more severe symptoms was related to the higher proportion of grass pea in staple food. Additionally, neuropathy could be easily prompted in high humidity and cold environment. A followed-up study 24 neuropathy sufferers for 23 years demonstrated that none of them recovered.¹¹

The research progress of grass pea in China

Since grass pea has a long history of cultivation as food for human consumption and feed for animal stock, it has been continuously researched and was found to be superior in many quality parameters such as strong resistance, nutritional value and nitrogen fixation. Current researches of grass pea in China are focused mainly on following six areas:

(1) *The extraction and detection of β -ODAP.* Zhang et al¹² have developed a method to analyze β -ODAP and its inactive isomer α -ODAP using 2,4-dinitrofluorobenzene (DNFB) for precolumn derivatization. Jiao et al¹³ have successfully separated β -ODAP in grass pea by thin layer chromatography. Zhao et al¹⁴ used pure water, 30% aqueous ethanol and 0.2 mol/L perchloric acid

to extract β -ODAP and found that 0.2 mol/L perchloric acid extraction combined with HPLC (high performance liquid chromatography) is an optimized method.

(2) *The screening of low toxin varieties.* Numerous works have been done to select low toxin varieties of grass pea. Bao et al¹⁵ have selected three varieties with higher yield and protein content and lower content of toxin from 65 varieties during three years. The mutagenic effect of heavy ion C^{6+} on the dry seed of grass pea has been studied by Wang et al,¹⁶ who discussed the suitable dose of C^{6+} for inducing low- or non-toxin genotypes of grass pea. Chinese researchers found that the processing of grass pea to vermicelli can dramatically decrease the content of β -ODAP to less than 1%.¹⁷ In addition, both microbial fermentation¹⁸ and two successive solid state fermentations¹⁹ could reduce the neurotoxin β -ODAP efficiently.

(3) *Investigation on the biological significance of β -ODAP.* β -ODAP was also discovered in the well-known traditional Chinese medicinal herb *Panax notoginseng*, in which it was named dencichine and used as a haemostatic.²⁰ Although β -ODAP can lead to neuropathy, it had no influence on animal reproduction. Feeding animals with enormous amount of grass pea for long term could not cause cancer or trigger tumour.²¹ Since the high level of β -ODAP and excellent drought resistance simultaneously exist in grass pea, Xing et al^{22,23} put forward an assumption: the function of ODAP may be similar to polyamine and proline under stress condition, because ODAP is not only a nitrogen compounds, but also a free amino acid and soluble in water, which can be regarded as osmotic regulation substance and an agent preventing dehydration to play an important role in nitrogen and energy metabolism. It provides a novel idea to study the drought resistance of grass pea.

(4) *The factors influencing β -ODAP content.* There were different levels of β -ODAP at different developmental stages, and the young leaves and mature seeds contain the highest level of β -ODAP.

Field studies have shown that β -ODAP levels in seeds were correlated positively with P, and negatively with Zn, N, Ca, K or Mg in the soil, and the level of β -ODAP in seedlings obviously reduced when infected by *Rhizobium*. It was also found that drought stress could increase the levels of β -ODAP in the seeds. In addition, β -ODAP accumulation could be related to low levels of endogenous H_2O_2 in the leaves under either field conditions or osmotic stress.²⁴

(5) *Studies on drought resistance.* Sun et al²⁵ treated *L. sativus* seedlings with PEG6000 to explore the physiological mechanism. Through a series of experiments, they suggested that the seedlings of *L. sativus* could alleviate its stress damage by the reduction of stomatal conductance/transpiration rate and the increased accumulation of proline, which might be a drought resistant mechanism. Jiang et al²⁶ also certified that the accumulation of osmoprotectants and the improvement of oxidation resistance resulted in the higher drought tolerance of grass pea compared to pea. Tian et al²⁷ have shown that Eu^{3+} induces abscisic acid early and promotes accumulation of ABA in grass pea under drought stress. In order to further explore the drought resistance mechanism of grass pea, Wu et al²⁸ established two-dimensional electrophoresis of leaf proteins of *L. sativus*, which lay the foundation for further proteomics research. Meanwhile, a recent finding²⁹ indicated that the peroxidase activity of grass pea becomes higher than in the control when treated with Cu^{2+} . Moreover, the level of reactive oxygen species (ROS) in grass pea increased significantly when it suffered from drought stress, and reduced glutathione (GSH) plays an essential role in ROS balancing, therefore it acts as an important molecule in the drought resistance of grass pea. Yang et al³⁰ developed fluorescence spectrophotometry for measurement of GSH in grass pea tissues, which contributed to the studies for resistance mechanisms of grass pea.

(6) *On other aspects:* Jiang et al³¹ found that exogenous hydrogen peroxide reversibly inhibited root gravitropism and induced horizontal curvature

of primary root during grass pea germination. Xu Shanshan³² explored the optimal condition to extract protein from *L. sativus*. Zhang et al³³ optimized the extracting conditions of grass pea starch, which combined immersion method with protein enzymolysis using Alkaline protease. Ecological studies on grass pea were mainly focused on the growth and development regularity, floral dynamics and breeding system of *L. quinquenervius*, and its community ecology.^{34,35,36}

Although grass pea has a number of unique features that make it attractive to the grower and consumer, there are still plenty of works that remain to be done: i) Most people know little about grass pea so that they buy and consume grass pea seedlings in their daily life unwittingly. Therefore dietary information, education, and communication on safe grass-pea preparations should be developed. ii) With regard to the screening of low-toxin grass pea species, molecular biological technique should be used to produce new grass pea genotypes which are safe for human consumption. iii) Meanwhile, grass pea is an efficient nitrogen fixer and is also a very intensively nodulated legume and studies on its nitrogen fixation process will provide new insights into its cultivation. iv) The role of ODAP in the resistance to various abiotic stresses and nitrogen fixation of grass pea needs to be further explored. v) Moreover, the medicinal value of grass pea is also worthy to be further investigated.

References

- ¹ Chen YZ, Li ZX, Lu FH, Bao XG, Liu SZ, Liu XC, Zhang GW, Li YR. Studies on the screening of low toxic species of *Lathyrus*, analysis of toxins and toxicology. Journal of Lanzhou University. 1992, 28(3): 93-98.
- ² Zhang BL, Mu CS, Wang Y, Wang YJ. Study on floral dynamic and breeding system of *Lathyrus quinquenervius*. Acta Prataculturae Sinica. 2006, 15(2): 68-73.
- ³ Luozhou JC. Wild leguminous grasses resources in Gannan and its evaluation. Pratacultural Science. 1996, 13(5): 59-61.
- ⁴ Wan GD. The eastern of Hexi Corridor leguminous forage grass resources and its assessment. Pratacultural Science. 2003, 20(7): 5-7.

- ⁵ Song LJ, Zhou SF, Li XF. The development and utilization of forage plants resources in Huder forestry bureau of Inner Mongolia. Inner Mongolia Forestry Investigation and Design. 2001, 24: 50-54.
- ⁶ Wang YF, Yang WX, Wang CX, Wu GL. The forage plant resource of legume family (Leguminosae) in Gansu. Pratacultural Science. 2006, 23(3): 12-16.
- ⁷ http://hzs9707.kwt.17888.com/ws-3000624-c0002_10-cn/news_25546.shtml.
- ⁸ Guo XQ. Variety of wild vegetable resource in Longdong region and strategies for development and exploitation. Gansu Agr. Sci. and Techn. 2005, 2: 29-30.
- ⁹ Liu YL, Kang YF, Li ZH, Liu HK. Screening and evaluating sprout characteristics of grasspea (*Lathyrus sativus* L.). Journal of Plant Genetic Resources. 2012, 13(4): 639-642.
- ¹⁰ Pan FX, Lu JW, Liu W, Geng MJ, Li XK, Cao WD. Study on characteristics of decomposing and nutrients releasing of three kinds of green manure crops. Plant Nutrition and Fertilizer Science. 2011, 17(1): 216-223.
- ¹¹ Guo JY, Yuan JM, Kang DX. Study on motor neuron injury caused by *Lathyrus sativus* poisoning. Chin J Neurol. 1997, 30(4): 223-226.
- ¹² Zhang HX, Liu MC, Jin XS, Luan F, Li ZX. Determination of β -N-oxalyl-L- α , β -diaminopropionic acid in *Lathyrus Sativus* by precolumn derivatization with 1-Fluoro-2,4-dinitrobenzene. Chinese Journal of Analytical Chemistry. 2006, 34(1): 100-102.
- ¹³ Jiao CJ, Yang LJ, Lei XY, Ma XF, Yuan ZY, Yang XP, Lang FY, Li ZX. Separation of ODAP in *Lathyrus sativus* by thin layer chromatography. Amino Acid and Biotic Resources. 2007, 29(4):76-80.
- ¹⁴ Zhao XX, He YG, Deng J, Fang CY, Meng YF. Determination of β -N-oxalyl- α , β -diaminopropionic acid in *Lathyrus sativus* by HPLC with different extraction methods. Food Research and Development. 2011, 32(9): 145-148.
- ¹⁵ Bao XG, Lu FH, Liu SZ, Shu QP. Sifting and cultivating utilization of low toxin varieties of *Lathyrus*. Pratacultural Science. 1995, 12(5): 48-54.
- ¹⁶ Wang CY, Yang HM, Wang YF, Wei ZQ, Liu YY, Wang GL. Mutation effect of C^{6+} heavy ion on seed of *Lathyrus sativus* L. Hereditas. 1993, 15(1): 28-31.
- ¹⁷ Gansu Academy of Agricultural Sciences, College of Chemistry and Chemical Engineering of Lanzhou University. Toxin analysis and detoxification methods of *Lathyrus sativus*. Journal of Lanzhou University. 1975, 2: 45-65.
- ¹⁸ Yao MM, Li ZX, Huang K, Zhao XX, Meng Y, Meng YF. Studies on degrading neurotoxin ODAP in *Lathyrus sativus* by microbial fermentation. Journal of Sichuan University (Natural Science Edition). 2007, 44(2): 339-442.
- ¹⁹ Zhao XX, Li ZX, Meng YF. Reduction Efficiency of the neurotoxin β -ODAP in *Lathyrus sativus* by two successive solid state fermentation. Food and Fermentation Industries. 2007, 33(6): 37-40.
- ²⁰ Xie GX, Qiu YP, Qiu MF, Gao XF, Liu YM, Jia W. Analysis of dencichine in *Panax notoginseng* by gas chromatography-mass spectrometry with ethyl chloroformate derivatization. Journal of Pharmaceutical and Biomedical Analysis. 2007, 43: 920-925.
- ²¹ Liu XC, Zhang GW, Li YR, Wang JX. Toxicological study on grass pea (*Lathyrus sativus* L.) and its toxic-component BOAA. Scientia Agricultura Sinica. 1989, 22(5): 86-93.
- ²² Xing GS, Cui KR, Li J, Wang YF, Li ZX. Water stress and accumulation of β -N-Oxalyl-L- α , β -diaminopropionic acid in grass pea (*Lathyrus sativus*). J. Agric. Food Chem.. 2001, 49: 216-220.
- ²³ Xing GS, Zhou GK, Li ZX, Cui KR. Studies of polyamine metabolism and β -N-oxalyl-L- α , β -diaminopropionic acid accumulation in grass pea (*Lathyrus sativus*) under water stress. Acta Botanica Sinica. 2000, 42(10): 1039-1044.
- ²⁴ Jiao CJ, Jiang JL, Li C, Ke LM, Cheng W, Li FM, Li ZX, Wang CY. β -ODAP accumulation could be related to low level of superoxide anion and hydrogen peroxide in *Lathyrus sativus* L. Food and Chemical Toxicology. 2011, 49: 556-562.
- ²⁵ Sun XY, Sun W, Li ZX, Li FM, Wang CY. Physiological response of *Lathyrus sativus* L. to drought stress. Bulletin of Botanical Research. 2008, 28(5): 589-593.
- ²⁶ Jiang JL, Su M, Chen YR, Gao N, Jiao CJ, Sun ZX, Li FM, Wang CY. Correlation of drought resistance in grass pea (*Lathyrus sativus*) with reactive oxygen species scavenging and osmotic adjustment. Biologia. 2013, 68(2): 231-240.
- ²⁷ Tian HE, Gao YS, Zeng FL, Li FM, Shan L. Effects of Eu^{3+} on amino acid and protein metabolisms in *Lathyrus sativus*. Acta Botanica Boreali-Occidentalia Sinica. 2005, 25(6):1171-1177.
- ²⁸ Wu QF, W CY, Wang XY, Li ZX, Wang YF. Establishment of two-dimension electrophoresis of leaf proteins of *Lathyrus sativus*. Acta Botanica Boreali-Occidentalia Sinica. 2006, 26(7): 1330-1336.
- ²⁹ Wang J, Wang TP, Shi JJ. Effect of copper stress on POD activity and isozymes of *Lathyrus sativus*. Journal of Anhui Agri. Sci. 2011, 39(7): 3801-3802.
- ³⁰ Yang LJ, Dong XN, Wang XF, Jiao CJ, Gu SY. Determination of reduced glutathione in *Lathyrus sativus*

L. by fluorometric method. *Science and Technology of Food Industry*. 2011, 32(4): 397-404.

³¹ Jiang JL, Su M, Wang LY, Jiao CJ, Sun ZX, Cheng W, Li FM, Wang CY. Exogenous hydrogen peroxide reversibly inhibits root gravitropism and induces horizontal curvature of primary root during grass pea germination. *Plant Physiology and Biochemistry*. 2012, 53: 84-93.

³² Xu SS. Extraction of protein from *Lathyrus sativus* and study on its functional characters. *Modern Food Science and Technology*. 2006, 22(3): 38-41.

³³ Zhang GQ, ZhuGe YX, Luo QG, Tian XL, Li YY, Chen TF. Optimization conditions of *Lathyrus sativus* protein extraction and starch preparation. *Grain Processing*, 2012, 37(5): 38-42.

³⁴ Song YT, Zhou DW, Yang Y. Study on seasonal dynamics of aboveground biomass of *Leymus chinensis* + *Lathyrus quinquenervius* community on Songnen grassland. *Chinese Journal of Grassland*. 2008, 30(3): 37-41.

³⁵ Zhang BL, Wang DS, Wang YJ, Mu CS. Comparative study on aboveground growth rule and biomass of *Lathyrus quinquenervius* in Songnen grassland. *Journal of Jilin Agricultural University*. 2006, 28(4): 379-382.

³⁶ Zhang BL, Mu CS, Wang Y, Wang YJ. Study on floral dynamic and breeding system of *Lathyrus quinquenervius*. *Acta Prataculturae Sinica*. 2006, 15(2): 68-73.

Z.X. Sun, H.Z. Hou, L.N. Zhou, W. Cheng, C.Y. Wang*
MOE Key Laboratory of Cell Activities and Stress Adaptations,
School of Life Sciences, Lanzhou University,
Lanzhou, 730000, China

Impact of cross-sectoral approach to addressing konzo in DRC

Introduction

From December 2009 to October 2011, Action Against Hunger (ACF-USA) implemented an intervention in the Bandundu province of the Democratic Republic of Congo (DRC) addressing factors underlying the konzo epidemic affecting the population of Kwango district. The 'Integrated Programme for the Eradication of Konzo in the Territory of Kwango in DRC' project was financed by the European Union (EU) Food Facility and aimed to eradicate the disease through a cross-

sectoral approach focused on nutrition education and training, dietary diversification, improved water access and agricultural processing. The strategy also aimed to address the high rates of malnutrition seen in konzo cases (25.8% global acute malnutrition (GAM) prevalence in konzo affected children less than 18 years old, 69.3% of GAM in konzo affected adults). A total of 22,000 households benefited from these activities. The project was implemented in 396 villages in the highly affected areas of Kahemba, Kajiji, Feshi and Panzi and to a lesser extent in Kenge, Boko, Popokabaka, Kasongo Lunda, Wamba Luadi and Kitenda across the Territory of Kwango. Due to limited resources, project activities were not carried out in all konzo locations, but rather in the most affected villages.

Konzo is a sudden epidemic spastic paraparesis (paralytic) disease which leads to a permanent paralysis of the affected person's lower limbs. It is a neurological ailment triggered by sustained dietary exposure to cyanide present in improperly processed cassava. Overall, vulnerability to konzo is heightened by the combination of low protein intake (associated with low dietary diversity), poor soil conditions (which favour the cultivation and consumption of bitter cassava varieties high in cyanide), and lack of sufficient water resources for thorough processing.

Methods

The impact study was conducted across the area targeted by the project. A stratified sampling approach was used, with six of eleven intervention health zones selected purposively and 40 of 395 intervention villages selected randomly. In each selected village, six beneficiary households were randomly selected to participate in household surveys (234 in total). Household surveys were supplemented with information from key informants and focus groups.

Findings

Knowledge and attitudes on konzo and nutrition

ACF employed a community outreach and mobilisation approach through the creation of community cells as a forum for discussion on konzo and nutrition. These served as launch pads for a broadly based educational campaign which extended to churches, schools, training of local health professionals, community volunteers and leaders, traditional authorities, etc. Changes in knowledge at endline compared to baseline suggest that community outreach and education activities were effective in challenging long held local beliefs on konzo and nutrition. At project baseline, 74% of sampled population attributed the disease to a metaphysical origin, while 88% correctly noted the food-related causes of konzo at endline. Similar results were found regarding knowledge, attitudes and practice on prevention strategies based on effective messaging that encouraged appropriate processing of cassava and inclusion of protein in diets through incorporation of maize flour into *fufu* preparation and legumes (pulses) in the diet.

Food stocks and dietary diversification

ACF introduced two improved food crop varieties, *niébé* (cowpea) varieties *Vita 7* and *Muyaya*, and sweet cassava varieties *TME119*, *Mwuazi*, *Nsasi*, *Disanka* and *Butamu* to support increased consumption of sulphur amino acids methionine and cysteine contained in cereal and leguminous foods and complement consumption of traditional cyanide-heavy bitter cassava varieties with varieties low in cyanide. Notable improvement was seen in overall food stocks and diversity of food items held by households over the life of the project. The positive trend in diversity and volume of household food stocks may be attributed to project impact, in particular IEC activities around balanced diets and food processing and preparation, as well as external factors such as climate, crop disease and seasonal fluctuations.

Cassava retting techniques and water access

ACF also supported the installation of village based mills to increase access to maize and cassava

milling services and improve the quality of the flour. Finally, ACF implemented a variety of hydraulic constructions: public retting tanks to process cassava, boreholes, springs, rainwater harvesting systems and piped distribution networks. Knowledge of community leaders and member households of community cells regarding cassava retting and drying techniques similarly improved. Community leaders correctly reporting optimal processing time increased from 60% to 99%. Member households showed similar improvement of knowledge but constraints around access to processing sites and water quality limited optimal practice. At endline, a majority of households indicated they were processing cassava in rivers or ponds and had largely abandoned utilization of home retting techniques that rely on prolonged use of the same water, saturated in acid and less effective in cyanide detoxification.

Konzo incidence

Kahemba health zone was found to harbour the highest number of confirmed cases at project baseline (1,639), and placed among the top three zones for incidence (2.08%) largely due to its density of population and associated risk factors. A surveillance system for screening and identification of konzo cases in Kahemba health zone was established by the local health structure in 2009, with annual caseload an estimated 1,300 individuals in 2009. MoH educational activities and ACF integrated activities on konzo were launched in early to mid 2010, with a marked decrease in cases recorded that year. A further reduction in caseload between 2010 and 2011 was noted during the critical months of June, July and August with 47 new cases recorded in 2011, an 84% reduction in annual incidence. The greatest reduction in new cases was observed among the under 5 years age group.




Figure 1: Comparison of number of konzo cases in Kahemba health zone in 2010 and 2011

Conclusion

ACF's multi-tiered community outreach and education strategy proved effective in the dissemination of information on a large scale. The community cell approach allowed for a deep, sustained and broad based appropriation of messages and activities around nutrition education and konzo that would not otherwise have been possible.




Figure 2: ACF program area in Kwango district, Bandundu province, DRC

Placing community members in leadership positions to carry out sensitisation allowed local taboos to be effectively mitigated through open discussion. This approach also permitted the affected population to control the educational process, encouraging better appropriation of

messages, knowledge transfer and behaviour change. Impacts achieved through the community outreach and education approach were reinforced by improved access to water, agricultural processing infrastructure and opportunities to diversify diets.

Muriel Calo

mcalo@actionagainsthunger.org

Muriel Calo is the Senior Food Security & Livelihoods Advisor for ACF-USA who provided technical support to the ACF-USA DRC mission.

Further reading:

The original version of this article appeared in the Emergency Nutrition Network (ENN)'s Field Exchange, issue 44 in December 2012.

For more information or to receive the full French language study report, contact: Muriel Calo, email: mcalo@actionagainsthunger.org

Special thanks go to the whole of the ACF project and study team and the communities who participated in the study. The funding support of the European Union (EU) Food Facility Programme is gratefully acknowledged.

Safety and toxicity profile of grass pea (*Khesari dal*): A reappraisal in the Indian scenario

The disease Neurotoxicity has gradually disappeared over the past three decades from India in particular and may soon become a part of history and even our views on grass pea (*Khesari dal*, *Lathyrus sativus*) toxicity have seen paradigm shifts. Neurotoxicity is now recognized as a disease that results only following excessive consumption of the pulse as a staple for prolonged periods especially under drought and famine situations wherein it is also the only survival food for more than 98% of the population. *Khesari dal* cultivation continues in several parts of India without any adverse effects reported during the last three decades. Recently some states in India have even revoked the ban on its sale. Some of these developments thus necessitate a reappraisal especially in the context of several recent

developments and also lessons learnt from the Ethiopian incidence.^{1,2} This brief review is to highlight some of the more important features that need to be considered in this regard.

Homoarginine (HA) the first unusual amino acid characterized from grass pea has over the years gone into the background as it had no toxic attributes. However, HA is now recognized as a normal metabolite in humans and its importance in human health is growing.^{3,4} HA is a better substrate for Nitric oxide synthase (NOS) compared to arginine and could be a source of sustained generation of NO.⁵ The health benefits of NO both for the cardiovascular and cerebral metabolism are now universally recognized. HA is also a weak inhibitor of arginase and may contribute to a raise in arginine levels.⁶ This is evidenced by the increased nitrate excretion in individuals on a grass pea diet. NO is a “master signaling molecule” and its role in cardiovascular physiology and health are very well established. A daily intake or supplementation with *khesari dal* may be of an added advantage due to its HA content and make it an unique and prized functional food as it could have a vasodilator effect and prevent prehypertension from progressing to full blown hypertension, a major risk factor for heart attacks and strokes. Recent clinical studies do show that higher HA levels are associated with lowered blood pressure and a higher survival rate from cardiovascular events. Some genetic information suggests that lowered HA levels due to the deficiency of arginine glycine amidinotransferase (AGAT) may be the link to the pathological basis of acute ischemic stroke.⁷

New research also suggests that HA may be important for skeletal health and its deficiency may be associated with high bone turnover and low bone mineral density (BMD)⁸. This property may be related to the weak inhibitory effect of HA on alkaline phosphatase⁹. HA is also an inhibitor of glycine uptake at the synapse (Author *unpublished results*) and this may benefit LTP, one of the major

cellular mechanisms that underlie [learning](#) and [memory](#). Unlike arginine, HA is not a substrate for pancreatic arginase and thus may be a better insulin secretagogue compared to arginine.¹⁰

Much of the research on grass pea toxicity has always centered on ODAP, and its acute toxicity is now well demonstrated. Primate model studies have had some success but only under intrathecal or lumbar routes of ODAP administration or oral feeding of large quantities of the seeds, seed extracts or fortified seed extracts and complex feeding regimens.^{11,12} In humans much of the ingested ODAP is metabolized (detoxified) and very little of it is excreted in the urine (less than 2% in 24 hrs) while in rats, chicks or monkeys at least 95% of ODAP is excreted unchanged which suggests that humans have a unique ability to metabolize (oxidation?) ODAP.¹³

Some recent studies show that ODAP is a nonapoptotic activator of PKC followed by several downstream events which includes stabilization of HIF and expression of VEGF and eNOS.^{14,15,16} Some of these findings provide fresh leads into possible therapeutic applications of both ODAP and HA and utilization of grass pea for human health. The discovery of ODAP (Dencichin) as a constituent of *Panax Notoginseng* the traditional Chinese medicinal herb is a new twist to the ODAP and grass pea saga.¹⁷ Several of the curative attributes of this herb are now ascribed to its Dencichin (ODAP) content. Some of the attributes of ODAP (Dencichin) are now being explored for potential therapeutic applications. Some therapeutic applications of ODAP/Dencichin that have been patented very recently are a) US patent for its application as a hemostatic agent¹⁸ b) China patent for the Fourth Military Medical University for its use in preparing a medicament for treating neurodegenerative diseases¹⁹ and c) China patent for its use in toothpastes to prevent oral ulcers (98% efficacy) and for dental hygiene.²⁰ More patents based on use of ODAP in combating Hypoxia, stroke and other applications are very

likely in the future.

All the surveys done during the last 60-70 years with grass pea consuming cohorts have only focused on the incidence of neurolathyrism and finally we have only realized that it is a simple case of excessive consumption of grass pea as a staple for many years that is the root cause. The time has now come to change this approach and what is now needed is to give up the conventional reflex hammer approach but identify the positive contributions and effects that grass pea can offer.

A whole new approach is now possible with the newer inputs on ODAP and HA. It seems appropriate at this stage to ponder if grass pea consumption as part of a cereal based diet is fit as a functional food until we are able to fix its daily dietary intake limits and this would be the way to move forward to restore its deserved place in human health and nutrition.

References:

- ¹ Getahun H, Lambein F, Vanhoorne M, Van der Stuyft P. Food aid cereals to reduce neurolathyrism related to grass pea preparations during famine. *Lancet* 2003;362:1808-10.
- ² Getahun H, Mekonnen A, TekleHaimanot R, Lambein F. Epidemic of neurolathyrism in Ethiopia. *Lancet* 1999; 354: 306-7.
- ³ Rao SLN. A look at the brighter facets of β -N-oxalyl-L- α,β -diaminopropionic acid, homoarginine and the grass pea. *Food chem toxicol* 2011;49:620-22.
- ⁴ Marz W, Meinitzer A, Drechsler C, Pilz S, Krane V, Marcus E, *et al.* Homoarginine, Cardiovascular risk and Mortality. *Circulation* 2010;122:967-75.
- ⁵ Jyothi P, Rao SLN. Sustained nitric oxide generation with L-Homoarginine. *Res Comm in Biochem cell & Mol Biol* 1999; 3: 223-232.
- ⁶ Hrabak A, Bajor T, Temesi A. Comparison of substrate and inhibitor specificity of arginase and nitric oxide (NO) synthase for arginine analogues and related compounds in murine and rat macrophages. *Biochem Biophys Res Commun.* 1994; 198: 206–212.
- ⁷ Davids.M, Ndika,J.D, Salomons G.S, Blom.H.J, Teerlink.T. Promiscuous activity of arginine: glycine amidinotransferase is responsible for the synthesis of the novel cardiovascular risk factor homoarginine. *FEBS Lett* 2012;586: 3656-57.
- ⁸ Pilz S, Meinitzer A, Tomaschitz A, Kienreich K, Fahrleitner-palmer,Drechsler C,*et.al* Homoarginine deficiency is associated with increased bone turnover. *Osteoporosis int* 2012;23:2731-32.
- ⁹ Lin CW, Fishman WH. L-Homoarginine. An organ specific uncompetitive inhibitor of human liver and bone alkaline phosphatases. *J Biol Chem* 1972; 247: 54-60.
- ¹⁰ Henningsson R,Lundquist. L: Arginine-induced insulin release is decreased and glucagon increased in parallel with islet NO production: *Am J Physiol* 1998;275: E500-6.
- ¹¹ Spencer PS, Roy DN, Ludolph A, Hugon J, Dwivedi MP, Schamburg HH. Lathyrism: evidence for role of the neuroexcitatory amino acid BOAA. *Lancet* 1986; 2 (8515): 1066-67.
- ¹² Kusama-Eguchi K, Yamazaki Y, Ueda T, Suda A,Hirayama Y,Ikegami F, *et.al* Hind-limb paraparesis in a rat model for neurolathyrism associated with apoptosis and an impaired vascular endothelial growth factor system in . the spinal cord. *J Comp Neurol* 2010; 518:982-42.
- ¹³ Pratap Rudra.M.P, Raghuvveer Singh.M, Junaid.MA, Jyothi.P, Rao SLN. Metabolism of dietary ODAP in humans may be responsible for the low incidence of Neurolathyrism . *Clin Biochem* 2004; 37:318-22.
- ¹⁴ Raghuvveer Singh M, Pratap Rudra.M, Rao SLN, Singh.SS. *in Vitro* activation of protein kinase C by β -N-Oxalyl-L- α,β -Diaminopropionic acid, the *Lathyrus sativus* neurotoxin . *Neurochem Res* 2004; 29:1343-48.
- ¹⁵ Ke Q, Costa M. Hypoxia inducible factor-1 (HIF-1). *Mol Pharmacol* 2006; 70:1469-80.
- ¹⁶ Nalini J, Sandepta B, Rajesh M, Ilavazhagan G, Rao SLN. Singh SS. β -N-Oxalyl-L- α,β -diaminopropionic acid regulates mitogen-activated protein kinase signaling by down-regulation of phosphatidylethanolamine-binding protein 1. *J Neurochem* 2011; 118: 176-86.
- ¹⁷ Zheng Y, Matsura Y, Han H. Isolation and structure elucidation of a new amino acid derivative from ginseng, *Acta.Pharm.Sin* 1996; 31:191-95.
- ¹⁸ Guihua L, Ping Chen, Quiu Sun, Sang Fong. Composition and methods for treating hemorrhagic conditions, US20110160307, 2011,
- ¹⁹ Fourth Military Med.Univ: Notoginseng factor in preparing medicine for neurodegenerative diseases: CN102579418A, 2012
- ²⁰ Kunming Zhenua Pharm Fact.Ltd., Toothpaste containing Dencichin and preparation method thereof, CN 102379827, 2012

S.L.N.RAO

Prof of Biochemistry, (Retd), Osmania University,

Hyderabad, I500007, India

Email:slnrao213@rediffmail.com

CCDN News will consider for publication short articles and letters (1-3 pages A 4 double spaced) in English. Because CCDN News is a newsletter, full-size original papers or reviews cannot be considered for publication. Material published in CCDN News may be freely reproduced, but always indicate that it comes from CCDN News. Please send all correspondence to the CCDNN Coordinator: Prof Fernand Lambein, Ghent University Belgium. <fernand.lambein@gmail.com>